

NEOVELA FOR DOGS

- Kills adult fleas, flea eggs and flea larvae in the environment;
- Treats, controls and prevents flea infestations;
- Controls flea allergy dermatitis;
- Prevents heartworm disease caused by *Dirofilaria immitis*;
- Treats and controls ear mites (*Otodectes cynotis*);
- Treats and controls sarcoptic mange (*Sarcoptes scabiei*);
- Treats biting lice (*Trichodectes canis*) and sucking lice (*Linognathus setosus*);
- Waterproof after 2 hours in dogs.

NEOVELA FOR CATS

- Kills adult fleas, flea eggs and flea larvae in the environment;
- Treats, controls and prevents flea infestations in cats;
- Controls flea allergy dermatitis in cats;
- Prevents heartworm disease caused by *Dirofilaria immitis* in cats;
- Treats and controls ear mites (*Otodectes cynotis*) in cats;
- Treats ear mites (*Psoroptes cuniculi*) in rabbits;
- Treats sarcoptic mange (*Sarcoptes scabiei*) in rabbits;
- Treats and controls intestinal hookworm (*Ancylostoma tubaeforme*) and roundworm (*Toxocara cati*) in cats;
- Treats biting lice (*Felicola subrostratus*) in cats;
- Treats cheyletiellosis (*Cheyletiella sp.*) in rabbits;
- Waterproof after 2 hours in cats.

NEOVELA FOR PUPPIES AND KITTENS

- Kills adult fleas, flea eggs and flea larvae in the environment;
- Treats, controls and prevents flea infestations in puppies and kittens;
- Controls flea allergy dermatitis in puppies and kittens;
- Prevents heartworm disease caused by *Dirofilaria immitis* in puppies and kittens;
- Treats and controls ear mites (*Otodectes cynotis*) in puppies and kittens;
- Treats ear mites (*Psoroptes cuniculi*) in small rabbits;
- Treats and controls sarcoptic mange (*Sarcoptes scabiei*) in puppies;
- Treats sarcoptic mange in small rabbits;
- Treats and controls intestinal hookworm (*Ancylostoma tubaeforme*) and roundworm (*Toxocara cati*) in kittens;
- Treats biting (*Trichodectes canis*) and sucking lice (*Linognathus setosus*) in puppies and biting lice (*Felicola subrostratus*) in kittens;
- Treats cheyletiellosis (*Cheyletiella sp.*) in small rabbits;
- Waterproof after 2 hours in puppies and kittens.

DIRECTIONS FOR USE

RESTRAINTS:

DO NOT USE in rabbits that may be used or processed for human consumption.

CONTRAINDICATIONS:

This product is contraindicated for use on cats and dogs less than 6 weeks of age. The product is contraindicated for use on rabbits less than 8 weeks of age.

HOW TO APPLY:

1. Remove required tube(s) from the blister package.
2. Hold tube upright, peel the ring lock off, then twist off the cap.
3. Sites of Application:



Cats
At the back of the neck



Dogs
between the shoulder blades

4. The animal should be standing for easy application. Remove any collar prior to application and replace when product is dry. Part the hair until the skin is visible. Place the tip of the tube through the pet's hair to the skin and squeeze the tube firmly to ensure the entire contents are applied directly onto the skin.

Do not apply to broken skin. Do not massage the application site. Do not apply when the animal hair coat is wet. Do not apply the product superficially to the hair coat and avoid contact with the treated area until dry. Avoid pet's eyes or mouth. Avoid contact between the product and fingers. Bathing or shampooing the animal two or more hours after treatment will not reduce the product efficacy for dogs and cats. In rabbits, the effect of bathing or shampooing on the product efficacy has not been assessed.

DOSAGE AND ADMINISTRATION

Animal & Bodyweight	Tube Dose	Number of Tube per Month
PUPPIES, KITTENS OR RABBITS UP TO 2.5 KG	0.25mL	1
CATS & RABBITS 2.6 - 7.5 KG	0.75mL	1
DOGS 2.6 - 5 KG	0.25mL	1
DOGS 5.1 - 10 KG	0.5mL	1
DOGS 10.1 - 20 KG	1.0mL	1
DOGS 20.1 - 40 KG	2.0mL	1
FOR CATS OVER 7.5 KG	Appropriate combination of tubes	
FOR DOGS OVER 40 KG		

Apply topically to the skin of dogs and cats once a month. For single treatment of rabbits. For external use only.
DO NOT GIVE PRODUCT BY MOUTH.

NEOVELA

FLEAS & HEARTWORM

120 mg/mL SELAMECTIN

FOR DOGS 2.6 - 5 KG	0.25mL/TUBE
FOR DOGS 5.1 - 10 KG	0.5mL/TUBE
FOR DOGS 10.1 - 20 KG	1.0mL/TUBE
FOR DOGS 20.1 - 40 KG	2.0mL/TUBE

FLEAS, HEARTWORM & WORMS

60 mg/mL SELAMECTIN

FOR PUPPIES AND KITTENS UP TO 2.5 KG	0.25mL/TUBE
FOR CATS 2.6 - 7.5 KG	0.75mL/TUBE



NEOVE®

NEOVE PHARMA AUSTRALIA PTY LTD
ACN 140 367 442
303c, Level 3, 276 Pitt Street
Sydney NSW 2000
Infoline: 1300 052 066
www.neovepharma.com.au

APVMA APPROVAL NO.:
85279/117223, 85280/117222,
85281/117221, 84376/117220
84378/117224, 84377/117225

FOR ANIMAL TREATMENT ONLY READ SAFETY DIRECTIONS

NEOVELA (SELAMECTIN)

Topical Parasiticide for Dogs, Cats,
Puppies, Kittens and Rabbits

GENERAL DIRECTIONS

**NOT TO BE USED FOR ANY PURPOSE,
OR IN ANY MANNER, CONTRARY TO
THIS LABEL UNLESS AUTHORISED
UNDER APPROPRIATE LEGISLATION.**

FLEA TREATMENT, CONTROL AND PREVENTION

If the animal is already infested with fleas when the first dose of NEOVELA is applied, then adult fleas on the animal are killed and no viable flea eggs are produced after the first application. This stops flea reproduction. An environmental infestation of fleas may persist for a short time after beginning treatment with NEOVELA because of the emergence of adult fleas from pre-existing pupae. However, significant reductions in flea infestations are to be expected after just one monthly treatment as flea larvae in the animal's surroundings are killed or prevented from developing to adults by treatment with NEOVELA.

For the prevention and lasting control of flea infestations, NEOVELA should be applied at monthly intervals throughout the flea season, starting one month before fleas become active. This ensures that fleas infesting the animal are killed and that no viable flea eggs are produced by these fleas. Flea larvae in the animal's surrounds are also killed by NEOVELA. This breaks the flea life cycle, controls and prevents flea infestations and improves the clinical signs associated with flea allergy dermatitis in most animals. The clinical signs in some animals may not respond to this treatment alone.

To obtain maximum flea control, it is recommended that all cats and dogs in the same environment are treated.

HEARTWORM PREVENTION

For the prevention of heartworm disease NEOVELA must be applied on a monthly basis. NEOVELA may be applied year-round or at least within one month after the animal's first exposure to mosquitoes and monthly thereafter until the end of the mosquito season. This dose must be given within one month after the last exposure to mosquitoes. If a dose is missed and a monthly interval between dosing is exceeded, then immediate application of NEOVELA and resumption of monthly dosing will minimise the opportunity for the development of adult heartworms. When replacing another heartworm preventive product in a heartworm disease prevention program, the first dose of NEOVELA must be given within a month of the last dose of the former medication. NEOVELA may be safely applied to heartworm infected animals, however, it is recommended, in accordance with good veterinary practice, that all animals 6 months of age be tested for existing heartworm infections before beginning medication with NEOVELA. NEOVELA is not effective against adult *D. immitis*; however it may decrease the number of circulating microfilariae.

EAR MITE TREATMENT

A single dose of NEOVELA was 100% effective in treating ear mites (*O. cynotis*) in laboratory studies with dogs and cats. In field studies, with a potential for reinfestation, a single dose of NEOVELA eliminated ear mites in 100% of cats, and two monthly doses of NEOVELA eliminated ear mites in 90% of dogs. A single dose of NEOVELA was also found to be 100% effective in treating ear mites (*Psooptes cuniculi*) in rabbits. Monthly use of NEOVELA in dogs and cats will treat any subsequent ear mite infestations. NEOVELA is suitable for a single treatment in rabbits. Cleansing of the infested ears is recommended to remove the debris.

SARCOPTIC MANGE TREATMENT FOR DOGS AND RABBITS

For the elimination of Sarcoptic mange

(*Sarcoptes scabiei*), NEOVELA should be applied once and is highly efficacious as a single dose. A second monthly dose of NEOVELA may be required for complete elimination of mites in some dogs. Monthly use of NEOVELA in dogs will treat any subsequent *Sarcoptes* mite infestations. NEOVELA is suitable for a single treatment in rabbits.

NEMATODE TREATMENT FOR CATS

For the treatment of intestinal hookworms (*A. tubaeforme*) and roundworms (*T. cati*), NEOVELA should be applied once as a single treatment. Monthly use of NEOVELA will control hookworm and roundworm infections.

LICE TREATMENT

A single dose of NEOVELA was 100% effective in treating biting lice in dogs (*Trichodectes canis*) and cats (*Felicola subrostratus*) with no lice found in laboratory studies at all assessments between 7 and 42 days post treatment. A single dose of NEOVELA eliminated sucking lice (*Linognathus setosus*) in 90% of treated dogs when assessed 28 days after treatment.

CHEYLETIELLA TREATMENT FOR RABBITS

For the treatment of Cheyletiella mites (*Cheyletiella* sp.) in rabbits, a single dose of NEOVELA was effective at eliminating mites by 5 weeks post-treatment.

ANIMAL SAFETY

DOGS NEOVELA has been tested with no adverse reactions in dogs of over 100 different pure and mixed breeds including Collies. No adverse effects were observed when NEOVELA was applied at 10 times the recommended dose and in pregnant and lactating females, breeding males and females and dogs six weeks of age and older. NEOVELA was also applied at 3 times the recommended dose to heartworm infected dogs, and no adverse effects were observed.

CATS NEOVELA has been tested with no serious adverse effects in cats of over 15

different pure and mixed breeds. No adverse effects were noted when NEOVELA was applied at 10 times the recommended dose and in pregnant and lactating females, breeding males and females and cats six weeks of age and older. NEOVELA was also applied at 4 times the recommended dose to heartworm infected cats, and no adverse effects were observed. Accidental oral administration of NEOVELA to cats may induce salivation or vomiting.

RABBITS NEOVELA has been shown to be safe when applied at up to 3 times the recommended dose. The safety of NEOVELA has not been assessed following accidental oral ingestion or in breeding rabbits of either sex including pregnant or lactating does.

Use of NEOVELA in dogs and cats has occasionally been associated with transient application site reactions such as alopecia, coat colour change and pruritus. In well-controlled clinical studies in dogs and cats, NEOVELA was used safely in animals receiving other frequently used veterinary products such as vaccines, anthelmintics, antiparasitics, antibiotics, steroids, collars, shampoos and dips.

SAFETY DIRECTIONS

Will irritate the eyes. Avoid contact with eyes. If product in eyes, wash it out immediately with water. Wash hands after use.

ADDITIONAL USER SAFETY

Flammable - Keep away from heat, sparks, open flames or other sources of ignition.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone 131 126.

DISPOSAL

Dispose of used tubes and packaging by wrapping in paper and placing in garbage.

STORAGE

Store below 30°C (room temperature) in a dry place in the original package.