

DIRECTIONS FOR USE:

By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain or distress when using this product other than as directed on the label. For external application only.

DOSAGE

For animals of this type	Use 1 tube from this pack content
Cats & Kittens less than 4 kg	0.4mL
Cats over 4 kg	0.8mL

1. Snap off ONE blister pack along the segment line

2. Use scissors to cut across the top bubble section
(Do not harm the tube inside).

3. Peel off the foil from the cut line and take out the tube.

4. Twist off the cap from the tube. Part the hair on the neck at the base of the skull until the skin is visible. Place the tip of the tube on the skin and squeeze the tube firmly to expel the entire contents directly on the skin. Do not get the product in your cat's eyes or allow your cat to ingest this product. The product is bitter tasting and salivation may occur for a short time if the cat licks the product immediately after treatment. Treatment at the base of the skull will minimize the chance for the cat to lick the product. Do not allow the product to run off. (On very rare occasions skin reactions such as hair loss, redness and itching may occur, however this is usually a single occurrence and hair will grow back uneventfully.)


5. Site of Application



Apply 1 TUBE
at the base of the skull

6. Record treatment date and repeat monthly or as directed.

7. The solvent in NEOVET may mark certain materials including plastics, leather, fabrics and finished surfaces. Allow application site to dry before permitting contact with such materials.



Segment Line

WARNING

KEEP OUT OF REACH OF CHILDREN
FOR ANIMAL TREATMENT ONLY

NEOVET

FOR CATS

Active Constituents: 100 g/L IMIDACLOPRID, 10 g/L MOXIDECTIN

For use on cats and kittens from 9 weeks of age. One application lasts 4 weeks.

FOR THE TREATMENT AND PREVENTION OF

- fleas (*Ctenocephalides spp.*); Larval flea stages are killed in the surroundings of NEOVET treated pets. NEOVET can be used as part of a treatment strategy for flea allergy dermatitis (FAD).



FOR THE CONTROL OF

- roundworms (*Toxocara cati*, *Toxascaris leonine*) including larval (L4) and immature adult stages
- hookworms (*Uncinaria stenocephala**), including larval (L4) and immature stages of all species;

FOR THE TREATMENT AND CONTROL OF

- lungworm (*Aelurostrongylus abstrusus*)
- ear mites (*Otodectes cynotis*)

*not clinically significant in New Zealand

Registered to :

NEOVE PHARMA AUSTRALIA PTY LTD

Level 3, 276 Pitt Street,
Sydney NSW 2000 Australia

Distributed by :

FORTIS

FORTIS VETS LIMITED

131 Hilton Highway, Washdyke
Timaru, 7910, New Zealand

Customer Information Line: 0800 800 981

GENERAL INSTRUCTIONS

NEOVET has been shown experimentally to kill adult and larval fleas within 20 minutes. Death can occur due to contact by absorption via the flea intersegmental membranes. Stops fleas from feeding in 3-5 minutes. Fleas are not required to bite to receive a lethal dose.

NEOVET can be used as part of a treatment strategy for flea allergy dermatitis (FAD). NEOVET breaks the FAD cycle by eliminating fleas which are the cause. Animals with severe FAD may require additional veterinary treatment.

It is recommended that all cats in the same environment should be protected by NEOVET FOR CATS. NEOVET may be used as the only product for flea control as it breaks the life cycle and kills fleas emerging from the environment to reinfest the cat. New fleas may continue to emerge from the environment for six weeks or longer (*depending on climate*) after treatment is initiated. More rapid resolution of heavy environmental infestations may be obtained by additional environmental control measures using a suitable registered product.

WORM CONTROL

NEOVET controls the major gastrointestinal nematodes of cats but does not control tapeworms. Cats can become reinfested with worms but regular use of NEOVET will remove them. If worm problems persist consult a veterinarian.

TREATMENT AND CONTROL OF EAR MITES

For the elimination of ear mites (*Otodectes cynotis*), NEOVET should be applied once and is highly efficacious as a single dose. Cats not responding to the initial treatment may respond to a second monthly treatment however this has not been established for NEOVET. Such non responding animals should be monitored and veterinary advice sought if treatment is not successful. Monthly application of NEOVET will treat any subsequent ear mite infestation. To maximise efficacy gentle cleaning of the ear canal is recommended.

TREATMENT AND CONTROL OF LUNGWORM

For the treatment and control of lungworm (*Aelurostrongylus abstrusus*), NEOVET is highly efficacious as a single dose. Monthly application of NEOVET will control any subsequent lungworm infections.

KITTENS

Kittens may be treated with NEOVET from 9 weeks of age. If kittens younger than 9 weeks require flea control they may be safely treated with Neove Imidacloprid Spot On for Cats. For control of roundworms and hookworms kittens require worming at 4, 6 and 8 weeks of age. When worming kittens prior to commencing NEOVET use a specific worming agent. Once treatment with NEOVET every month is commenced worms will be continuously controlled.

PRECAUTIONS

Use with caution in sick, debilitated or underweight animals. Take care to avoid the product coming into contact with the eyes or mouth of the animal. Ingestion by the recipient or in-contact animals should be prevented. Transient salivation, vomiting and pupil dilation may follow accidental oral ingestion.

PREGNANCY AND LACTATION

No primary embryo-toxic, teratogenic or reproductive toxic effects have been observed with imidacloprid or moxidectin in laboratory species. Evidence suggests that no adverse effects are to be expected in pregnant and lactating cats. However, the safety of NEOVET has not been established during pregnancy and lactation.

READ LABEL BEFORE USE

WARNING May be harmful if inhaled, swallowed or absorbed through the skin. May cause eye irritation. May cause an allergic skin reaction. Moxidectin may have effects on the nervous system and on or lactation. Imidacloprid may have effects on the liver.

HANDLING PRECAUTIONS

Avoid inhalation and contact with skin and eyes. When handling wear eye protection and protective gloves. Do not eat, drink or smoke while using. Wash hands and exposed skin before meals and after use.

FIRST AID

If swallowed do NOT induce vomiting. For advice contact the National Poisons Centre 0800 POISON (0800 764766) or a doctor immediately. If skin or hair contact occurs remove contaminated clothing and flush skin and hair with running water. If splashed in eyes wash out immediately with water.

DISPOSAL

Dispose of used tubes by wrapping in paper and placing in garbage. Also contains: Benzyl Alcohol >50% w/w.

STORAGE

Store below 30°C.
(Room Temperature)

Registered pursuant to the ACVM Act 1997, No. A011604
See www.foodsafety.govt.nz for registration conditions.